

**CLAIMS:**

Subst  
C<sup>1</sup>  
5

1. A method of producing a library of mutant nucleic acid molecules comprising:

(a) obtaining a template nucleic acid;

(b) preparing a first oligonucleotide corresponding to a first desired mutation within said template nucleic acid;

(c) preparing a second oligonucleotide corresponding to a second desired mutation within said template nucleic acid;

(d) mixing the oligonucleotides prepared in said steps (b) and (c) so as to

10 hybridize said oligonucleotides to said template nucleic acid;

(e) subjecting the mixture of step (d) to the linear cyclic amplification reaction to produce a library of mutant template nucleic acids.

15 2. The method according to claim 1, wherein said oligonucleotides in said steps (b) and (c) are discontiguous.

3. The method according to claim 1, wherein said step first and second oligonucleotides are present in less than saturation concentration.

20 4. The method according to claim 1, wherein the mixture of said step (d) further comprises non-mutagenic oligonucleotides corresponding to either or both of said first and second oligonucleotides.

25 5. The method according to claim 1, wherein said template nucleic acid corresponds to a desired protein product.

6. The method according to claim 4, wherein said protein product comprises an enzyme, hormone, vaccine, peptide therapeutic or antibody.

30 7. The method according to claim 4, further comprising the steps of:

(f) transforming said mutant template nucleic acids from said library into a competent host cell;

(g) expressing protein corresponding to said mutant nucleic acids in said host cell;

35 (h) screening said expressed proteins for desired characteristics.

add  
C<sup>1</sup>